



<i>Phase</i>	<i>Material Name</i>	<i>US INCI</i>	<i>Supplier</i>	<i>% Material</i>
A	Deionized Water	Water	N.A.	54.69
A	Optigel-WX	Bentonite Xanthan Gum	BYK	2.00
A	1,2-Propanediol	Propylene Glycol	VWR	2.50
A	Citric Acid (10%)	Water Citric Acid	N.A.	0.01
B	Emulium Delta MB	Cetyl Alcohol Glyceryl Stearate PEG-75 Stearate Ceteth-20 Steareth-20	Gattefossé	5.00
B	Lipocire A Pellets	C10-18 Triglycerides	Gattefossé	2.00
B	Compritol 888 CG Pellets	Glyceryl Behenate	Gattefossé	2.00
B	Cocoate BG MB	Butylene Glycol Cocoate	Gattefossé	4.00
B	Tocomix L50-IP	Tocopherol Helianthus Annuus (Sunflower) Seed Oil	Jan Dekker	0.30
B	KahlWax 1899 Micro	Microcrystalline Wax	KahlWax	2.00
C	SynCrystal Almond	Synthetic Fluorphlogopite Titanium Dioxide Iron Oxides Tin Oxide	Eckart	12.00
C	SynCrystal Soft Black	Synthetic Fluorphlogopite Iron Oxides	Eckart	4.00
D	Euxyl PE9010	Phenoxyethanol Ethylhexylglycerin	Schuelke	1.00
D	MultiMoist CLR™	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	3.00
D	Follicusan™ DP	Water Alcohol Denat. Panthenyl Ethyl Ether Milk Protein Lactose Inositol Acetyl Cysteine Acetyl Methionine Sodium Citrate Citric Acid	CLR	5.00
D	Vitamin F forte	Linoleic Acid Linolenic Acid	CLR	0.50
				100.00

Operating Instructions

Disperse OPTIGEL-WX into water and stir for 10 minutes. Adjust pH value to 6.5 with Citric Acid and add Propylene Glycol. Heat up A and B to 80°C separately. Add B to A and cool down while stirring. Add C at 70°C and stir cold. Add D in the given order below 40°C. Mix until uniform and pour into an appropriate container.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com

PROJECT NAME : MultiMoist CLR™



FORMULA NAME : 3in1 Lash & Eyebrow Treatment

TRIAL CODE : 385.021.13

Directions for use:

For lashes- apply at base of lashes and brush upwards to volumize and lengthen. Layer as desired.

As eyebrow treatment- apply the brush slightly above your eyebrow and make sure the brush goes according to your natural growth.

As mascara- curl the lashes well and apply mascara starting from the roots of the top lashes, moving the wand first side to side then upwards toward the end of the lashes.

*developed & produced by Eckart; www.eckart.net

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com